

Explicit CP violation in a MSSM with an extra $U(1)'$

S.W. Ham⁽¹⁾, E.J. Yoo⁽²⁾, and S.K. Oh^(1,2)

⁽¹⁾ *Center for High Energy Physics, Kyungpook National University
Daegu 702-701, Korea*

⁽²⁾ *Department of Physics, Konkuk University, Seoul 143-701, Korea*

Abstract

We study that a minimal supersymmetric standard model with an extra $U(1)'$ gauge symmetry may accommodate the explicit CP violation at the one-loop level through radiative corrections. This model is CP conserving at the tree level and cannot realize the spontaneous CP violation for a wide parameter space at the one-loop level. In explicit CP violation scenario, we calculate the Higgs boson masses and the magnitude of the scalar-pseudoscalar mixings in this model at the one-loop level by taking into account the contributions of top quarks, bottom quarks, exotic quarks, and their superpartners. In particular, we investigate how the exotic quarks and squarks would affect the scalar-pseudoscalar mixings. It is observed that the size of the mixing between the heaviest scalar and pseudoscalar Higgs bosons is changed up to 20 % by a complex phase originated from the exotic quark sector of this model.

I. INTRODUCTION

Possessing two Higgs doublets is one of the characteristics of the minimal supersymmetric standard model (MSSM) in order to give masses to the up-quark sector and the down-quark sector separately [1]. This property enables in principle CP violation to occur in the Higgs sector of the MSSM through the mixing between the CP even and the CP odd states [2]. The mechanism of CP violation in the MSSM has been studied by many authors [3-12]. It is found that at the tree level, neither explicit nor spontaneous CP violation is possible in the Higgs sector of the MSSM, because any complex phases in it can always be eliminated by rotating the Higgs fields. Even at the one-loop level, spontaneous CP violation is disfavored because it requires a very light neutral Higgs boson, which has already been ruled out by experiments [3,4]. On the other hand, explicit CP violation is viable in the MSSM at the one-loop level by virtue of the radiative corrections due to the loops of relevant particles, such as quarks and squarks [5-12]. The radiative corrections by these particles yield the mixing between the CP even and the CP odd neutral Higgs bosons. Thus, it is possible to achieve the explicit CP violation in the radiatively corrected Higgs sector of the MSSM.

However, a drawback of the MSSM is that it contains a Higgs mixing term $\mu H_1 H_2$, where μ is a parameter having mass dimension, and H_i ($i = 1, 2$) are Higgs doublets. It is known that this μ parameter causes a hierarchy problem with respect to the symmetry breaking scale [13]. A natural framework to solve this problem is to generate the μ parameter via the vacuum expectation value (VEV) of a Higgs singlet [14]. A possibility for the framework is to extend simply the Higgs sector of the MSSM by introducing a Higgs singlet, like the next-to-minimal supersymmetric standard model, where the μ parameter is practically replaced by a dimensionless parameter times the VEV of the Higgs singlet [14].

A more plausible possibility to solve the μ problem is to extend not only the Higgs sector but also the gauge symmetry of the MSSM by introducing a nonanomalous $U(1)'$ symmetry broken at the TeV scale [15]. Extending the gauge symmetry to include an additional $U(1)'$ symmetry is widely considered in various theoretical models, such as string models or GUT models. The MSSM with an extra $U(1)'$ symmetry can actually forbid the $\mu H_1 H_2$ term but allow $\lambda S H_1 H_2$, by assigning appropriate $U(1)'$ charges, where S is the Higgs singlet and λ is a dimensionless parameter. After S develops a VEV of the electroweak scale, this λ term would effectively generate the μ term at the electroweak scale [16].

In the literature, a number of studies have been performed on the MSSM with an extra $U(1)'$ symmetry [17-20]. Several authors have calculated the Higgs boson masses from the tree-level Higgs potential of this model [17] as well as from the radiatively corrected Higgs potential [18,19]. It is found that, at the tree level, the lightest scalar Higgs boson in this model can be heavier than Z boson. That is, the upper bound on the mass of the lightest scalar Higgs boson in this model is larger than the one in the MSSM without an extra $U(1)'$ symmetry. Moreover, it is also found that this tree-level mass of the lightest scalar Higgs boson in this model can be significantly affected by radiative corrections.

Recently, this model has been further investigated within the context of the supersym-

metric CP problem [20], as it can have explicit CP violating phases in the Higgs sector. At the tree level, like the MSSM, this model has no CP mixing between the CP even and the CP odd states in the Higgs sector because the single CP phase arising from the tree-level Higgs potential can always be eliminated by rotating the Higgs fields. In Ref. [20], the one-loop contribution from the top and stop quark loops are considered in the explicit CP violation scenario. It is observed that the lightest neutral Higgs boson remains essentially CP even for a wide parameter space, with a fixed value of $\tan\beta = 1$, where $\tan\beta = v_2/v_1$ is the ratio of the VEVs of two Higgs doublets. Meanwhile, large CP mixtures for the other heavier Higgs bosons may be realized as the size of the effective μ terms becomes large.

In this paper, motivated by the results of Ref. [20], we would like to study this model in some detail in the explicit CP violation scenario. The radiative CP mixing may be generated by means of the complex phases coming from the one-loop effective potential due to top quarks, bottom quarks, the exotic quarks, and their superparticles, where the exotic quarks and squarks with electric charges $\pm 1/3$ are introduced into this model in order to cancel gauge anomaly [21]. In particular, we are interested in the contributions of exotic quark sector in this model. At the one-loop level, the exotic quarks and squarks are found to play some recognizable role in the scalar-pseudoscalar mixings. We find that the relevant CP phase from the exotic quark sector may change the size of the CP mixing between the heaviest scalar and the pseudoscalar Higgs bosons by up to 20 %. Also we find that this model yields a negative or extremely small axion mass, as well as other unacceptable predictions, at the one-loop level for full parameter space in the spontaneous CP violation scenario. Thus, we note that the present model is impossible to realize spontaneous CP violation at the one-loop level.

II. HIGGS SECTOR

GUT, such as the string-inspired E_6 model, might be a natural motivation to enlarge the gauge symmetry in order to accommodate an extra $U(1)'$ symmetry into the MSSM. The E_6 gauge group may be decomposed into

$$E_6 \supset SU(10) \times U(1)_\psi \supset SU(5) \times U(1)_\chi \times U(1)_\psi, \quad (1)$$

where $SU(5)$ is further broken down to the Standard Model (SM) gauge group, $SU(3)_C \times SU(2)_L \times U(1)_Y$. At the electroweak scale, the desired extra $U(1)'$ symmetry may be given as an orthogonal linear combination of $U(1)_\chi$ and $U(1)_\psi$ as

$$Q' = \cos\theta_E Q_\chi + \sin\theta_E Q_\psi, \quad (2)$$

where Q' , Q_χ , and Q_ψ are the $U(1)'$, $U(1)_\chi$, and $U(1)_\psi$ charges, respectively. Sometimes, for four different values of the angle θ_E , the four particular combinations of the $U(1)_\chi$ and $U(1)_\psi$ are called as follows: the χ -model for $\theta_E = 0$, the ψ -model for $\theta_E = \pi/2$, the η -model for $\theta_E = \tan^{-1}(-\sqrt{5/3})$, and the ν -model for $\theta_E = \tan^{-1}\sqrt{15}$. The gauge group we consider is thus $G = SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)'$.

The Higgs sector of the MSSM with an extra $U(1)'$ symmetry consists of two Higgs doublets $H_1 = (H_1^0, H_1^-)$ and $H_2 = (H_2^+, H_2^0)$, and a neutral Higgs singlet S . This model possesses an extra pair of $SU(2)$ singlet quarks, D_L and $\bar{D}_R$, with electric charges $-1/3$ and $+1/3$, respectively, introduced by the requirement of the gauge anomaly cancellation. For the fermion matter fields in this model, we take into account only the third generation of quarks, besides the exotic quarks. Then, the superpotential for the model we consider may be expressed as

$$W \approx \lambda S H_1 H_2 + h_t Q H_2 t_R^c + h_b Q H_1 b_R^c + k S D_L \bar{D}_R, \quad (3)$$

where Q is the left-handed quark doublet, t_R^c and b_R^c are the charge conjugate of the right-handed top quark and bottom quark, respectively, with h_t and h_b being their respective Yukawa coupling coefficients, and $H_1 H_2 = H_1^0 H_2^0 - H_1^- H_2^+$. All the coupling coefficients are dimensionless. This superpotential is symmetric under $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)'$, and the $U(1)'$ symmetry is broken down by the VEV of S .

The Higgs potential of this model at the tree level may expressed as a sum of the F -term, the D -term, and the soft breaking term, that is,

$$V_0 = V_F + V_D + V_S, \quad (4)$$

where

$$\begin{aligned} V_F &= |\lambda|^2 [(|H_1|^2 + |H_2|^2)|S|^2 + |H_1 H_2|^2], \\ V_D &= \frac{g_2^2}{8} (H_1^\dagger \vec{\sigma} H_1 + H_2^\dagger \vec{\sigma} H_2)^2 + \frac{g_1^2}{8} (|H_1|^2 - |H_2|^2)^2 \\ &\quad + \frac{g_1'^2}{2} (\tilde{Q}_1 |H_1|^2 + \tilde{Q}_2 |H_2|^2 + \tilde{Q}_3 |S|^2)^2, \\ V_S &= m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_3^2 |S|^2 - [\lambda A_\lambda (H_1 H_2) S + \text{H.c.}], \end{aligned} \quad (5)$$

where m_i^2 ($i = 1, 2, 3$) are soft masses, g_2 , g_1 , and g_1' are respectively the coupling coefficients of $SU(2)$, $U(1)$, and $U(1)'$; $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the Pauli matrices; $\tilde{Q}_1$, $\tilde{Q}_2$, and $\tilde{Q}_3$ are respectively the $U(1)'$ charges of H_1 , H_2 , and S , satisfying the condition of $\tilde{Q}_1 + \tilde{Q}_2 + \tilde{Q}_3 = 0$ to obey the gauge invariance of the superpotential under $U(1)'$.

At the electroweak scale, the three neutral Higgs fields develop VEVs as $\langle H_1^0 \rangle = v_1$, $\langle H_2^0 \rangle = v_2$, and $\langle S \rangle = s e^{i\phi_s}$, where ϕ_s is the relative phase between $H_1 H_2$ and S . In general, λA_λ in the soft breaking term may be complex, like $\lambda A_\lambda e^\phi$. However, it is always possible to make the phase ϕ and the phase ϕ_s cancel out each other by redefining the Higgs singlet S , and thus the tree-level Higgs potential can be made free of any complex phase. Therefore, the CP symmetry is not violated in this model at the tree level.

Now let us consider the one-loop radiative corrections to the tree-level Higgs potential. In supersymmetric models, the incomplete cancellation between ordinary particles and their superpartners yield the one-loop corrections to the tree-level Higgs boson masses. Generally, the most dominant part of the one-loop corrections to the tree-level Higgs potential come primarily from the top and stop quark loops. By considering the top quark sector only, it has been observed that explicit CP violation in this model is viable through the radiatively corrected Higgs potential [20]. For large $\tan \beta$, the contribution of the bottom and sbottom quark loops can also be significantly large. We would like to

consider in this paper the one-loop contributions from both the top and bottom quark sector, as well as from the exotic quark sector.

To start with, we calculate the tree-level masses of the relevant particles in order to evaluate the radiative corrections to the neutral Higgs sector. The fermion matter fields obtain their masses after the electroweak symmetry breaking as $m_t^2 = (h_t v_2)^2$ for top quark, $m_b^2 = (h_b v_1)^2$ for bottom quark, and $m_k^2 = (k s)^2$ for the exotic quark. The tree-level masses of their superpartners are given as

$$\begin{aligned} m_{\tilde{t}_1, \tilde{t}_2}^2 &= m_T^2 + m_t^2 \mp m_t \sqrt{A_t^2 + \lambda^2 s^2 \cot^2 \beta - 2\lambda A_t s \cot \beta \cos \phi_t} , \\ m_{\tilde{b}_1, \tilde{b}_2}^2 &= m_B^2 + m_b^2 \mp m_b \sqrt{A_b^2 + \lambda^2 s^2 \tan^2 \beta - 2\lambda A_b s \tan \beta \cos \phi_b} , \\ m_{\tilde{k}_1, \tilde{k}_2}^2 &= m_K^2 + m_k^2 \mp m_k \sqrt{A_k^2 + \lambda^2 v^4 \sin^2 2\beta / (4s^2) - \lambda A_k v^2 \sin 2\beta \cos \phi_k / s} , \end{aligned} \quad (6)$$

where $\tan \beta = v_2/v_1$, $v^2 = v_1^2 + v_2^2$. Further, m_T , m_B , and m_K are the soft SUSY breaking masses respectively for stop quarks, sbottom quarks, and the exotic squarks, and likewise A_t , A_b , and A_k are respectively their trilinear soft SUSY breaking parameters of the mass dimension. One can notice that there are three phases ϕ_t , ϕ_b , and ϕ_k in the above expressions. These complex phases are determined by the generally complex A_q ($q = t, b, k$) and the phase of s , the VEV of the Higgs singlet. We note that the D -terms do not contribute to the squark masses.

The full Higgs potential at the one-loop level may be written as

$$V = V_0 + V_1 ,$$

where V_1 is the radiative corrections due to the relevant particles and their superpartners. According to the effective potential method [22], V_1 is given as

$$V_1 = \sum_l \frac{n_l \mathcal{M}_l^4}{64\pi^2} \left[\log \frac{\mathcal{M}_l^2}{\Lambda^2} - \frac{3}{2} \right] , \quad (7)$$

where Λ is the renormalization scale in the modified minimal subtraction scheme, and the subscript l stands for various participating particles: $t, b, k, \tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2, \tilde{k}_1$ and $\tilde{k}_2$. The degrees of freedom for these particles including the sign convention are $n_t = n_b = n_k = -12$ and $n_{\tilde{t}_i} = n_{\tilde{b}_i} = n_{\tilde{k}_i} = 6$ ($i = 1, 2$), since in the above formula enter fermions with a negative sign while bosons with a positive sign.

If CP violation takes place in the Higgs sector, the neutral Higgs bosons would not have definite CP parity, hence mixings among them. In case of explicit CP violation, the non-trivial tadpole minimum condition with respect to the pseudoscalar component of the Higgs field is given as

$$\begin{aligned} 0 &= A_\lambda \sin \phi_0 - \frac{3m_t^2 A_t \sin \phi_t}{16\pi^2 v^2 \sin^2 \beta} f(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2) - \frac{3m_b^2 A_b \sin \phi_b}{16\pi^2 v^2 \cos^2 \beta} f(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2) \\ &\quad - \frac{3m_k^2 A_k \sin \phi_k}{16\pi^2 s^2} f(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2) , \end{aligned} \quad (8)$$

where the dimensionless function f arising from radiative corrections is defined as

$$f(m_x^2, m_y^2) = \frac{1}{(m_y^2 - m_x^2)} \left[m_x^2 \log \frac{m_x^2}{\Lambda^2} - m_y^2 \log \frac{m_y^2}{\Lambda^2} \right] + 1 ,$$

and the phase ϕ_0 is given by $\phi_0 = \phi_s + \phi$, with ϕ being the phase of λA_λ in V_0 and ϕ_s being the phase of s . Note that ϕ_0 is generally nonzero at the one-loop level whereas at the tree level it can be made zero by rotating the Higgs singlet. In the above tadpole minimum condition, the first term comes from the tree-level Higgs potential, and the remaining three terms come respectively from the top squark, bottom squark, and exotic squark contributions.

At the tree level, the Higgs sector in the MSSM with an extra $U(1)'$ has ten real degrees of freedom, which are decomposed by two neutral Goldstone bosons, a pair of charged Goldstone bosons, four neutral Higgs bosons and a pair of charged Higgs bosons. After the electroweak symmetry breaking, the two neutral Goldstone bosons and a pair of charged Goldstone bosons will eventually be absorbed into the longitudinal component of Z , Z' , and W gauge bosons. Since the CP symmetry is conserved in the Higgs sector at the tree level, the four neutral Higgs bosons can be divided into three scalar Higgs bosons and one pseudoscalar Higgs boson, according to the CP parity.

The squared mass matrix M of the four neutral Higgs bosons is given as a symmetric 4×4 matrix that is obtained by the second derivatives of the Higgs potential with respect to the four Higgs fields, in the (h_1, h_2, h_3, h_4) basis. At the tree level, M is given by V^0 as

$$M = M^0 = \begin{pmatrix} M_{11}^0 & M_{12}^0 & M_{13}^0 & 0 \\ M_{21}^0 & M_{22}^0 & M_{23}^0 & 0 \\ M_{31}^0 & M_{32}^0 & M_{33}^0 & 0 \\ 0 & 0 & 0 & m_A^2 \end{pmatrix}, \quad (9)$$

where the tree-level squared mass of the pseudoscalar Higgs boson is given as

$$m_A^2 = 2\lambda A_\lambda v \frac{\cos \phi_0}{\sin 2\alpha}$$

with

$$\tan \alpha = \frac{v}{2s} \sin 2\beta$$

standing for the splitting between an extra $U(1)'$ symmetry breaking scale and the electroweak scale. Note that M^0 is divided into two blocks of the upper-left 3×3 submatrix and the single element $M_{44}^0 = m_A^2$, corresponding to the scalar part and the pseudoscalar part. There is no scalar-pseudoscalar mixing in M^0 , since M_{i4}^0 ($i = 1 - 3$) are zero, and hence no CP violation. Explicit calculations yield

$$\begin{aligned} M_{11}^0 &= m_Z^2 \cos^2 \beta + 2g_1'^2 \tilde{Q}_1^2 v^2 \cos^2 \beta + m_A^2 \sin^2 \beta \cos^2 \alpha, \\ M_{22}^0 &= m_Z^2 \sin^2 \beta + 2g_1'^2 \tilde{Q}_2^2 v^2 \sin^2 \beta + m_A^2 \cos^2 \beta \cos^2 \alpha, \\ M_{33}^0 &= 2g_1'^2 \tilde{Q}_3^2 s^2 + A^2 \sin^2 \alpha, \\ M_{12}^0 &= g_1'^2 \tilde{Q}_1 \tilde{Q}_2 v^2 \sin 2\beta + (\lambda^2 v^2 - m_Z^2/2) \sin 2\beta - m_A^2 \cos \beta \sin \beta \cos^2 \alpha, \\ M_{13}^0 &= 2g_1'^2 \tilde{Q}_1 \tilde{Q}_3 v s \cos \beta + 2\lambda^2 v s \cos \beta - m_A^2 \sin \beta \cos \alpha \sin \alpha, \\ M_{23}^0 &= 2g_1'^2 \tilde{Q}_2 \tilde{Q}_3 v s \sin \beta + 2\lambda^2 v s \sin \beta - m_A^2 \cos \beta \cos \alpha \sin \alpha, \end{aligned} \quad (10)$$

At the one-loop level, the squared mass matrix may be decomposed as

$$M = \bar{M}^0 + M^1$$

where $\bar{M}^0$ has exactly the same appearance as M^0 but m_A^2 in M^0 should be replaced by $\bar{m}_A^2$ in $\bar{M}^0$, where $\bar{m}_A^2$ is given as

$$\bar{m}_A^2 = \frac{2\lambda v}{\sin 2\alpha} \left[A_\lambda \cos \phi_0 - \frac{3m_t^2 A_t \cos \phi_t}{16\pi^2 v^2 \sin^2 \beta} f(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2) - \frac{3m_b^2 A_b \cos \phi_b}{16\pi^2 v^2 \cos^2 \beta} f(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2) - \frac{3m_k^2 A_k \cos \phi_k}{16\pi^2 s^2} f(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2) \right], \quad (11)$$

One may notice that in the expression for $\bar{m}_A^2$, the first term comes from the tree-level Higgs potential and the remanning three terms comes from the contributions of the squarks. Therefore, $\bar{M}^0$ contains not only the tree-level results but also the contributions from radiative corrections.

Now, let us calculate M^1 , which is obtained from V_1 . We may conveniently decompose M^1 as

$$M^1 = M^t + M^b + M^k$$

where symmetric matrices M^t , M^b , and M^k are obtained respectively from the top quark sector, the bottom quark sector, and the exotic quark sector contributions to V_1 , after imposing the tadpole minimum condition.

Somewhat lengthy calculations yield the matrix elements of M^t as

$$\begin{aligned} M_{11}^t &= \frac{3m_t^4 \lambda^2 s^2 \Delta_{\tilde{t}_1}^2}{8\pi^2 v^2 \sin^2 \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2}, \\ M_{22}^t &= \frac{3m_t^4 A_t^2 \Delta_{\tilde{t}_2}^2}{8\pi^2 v^2 \sin^2 \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} + \frac{3m_t^4 A_t \Delta_{\tilde{t}_2}}{4\pi^2 v^2 \sin^2 \beta} \frac{\log(m_{\tilde{t}_2}^2/m_{\tilde{t}_1}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)} \\ &\quad + \frac{3m_t^4}{8\pi^2 v^2 \sin^2 \beta} \log\left(\frac{m_{\tilde{t}_1}^2 m_{\tilde{t}_2}^2}{m_t^4}\right), \\ M_{33}^t &= \frac{3m_t^4 \lambda^2 \Delta_{\tilde{t}_1}^2}{8\pi^2 \tan^2 \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2}, \\ M_{44}^t &= \frac{3m_t^4 \lambda^2 A_t^2 s^2 \sin^2 \phi_t}{8\pi^2 v^2 \sin^4 \beta \cos^2 \alpha} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2}, \\ M_{12}^t &= -\frac{3m_t^4 \lambda A_t s \Delta_{\tilde{t}_1} \Delta_{\tilde{t}_2}}{8\pi^2 v^2 \sin^2 \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} - \frac{3m_t^4 \lambda s \Delta_{\tilde{t}_1}}{8\pi^2 v^2 \sin^2 \beta} \frac{\log(m_{\tilde{t}_2}^2/m_{\tilde{t}_1}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)} \\ M_{13}^t &= \frac{3m_t^4 \lambda^2 s \Delta_{\tilde{t}_1}^2}{8\pi^2 v \sin \beta \tan \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} - \frac{3m_t^4 \lambda^2 s \cos \beta}{8\pi^2 v \sin^2 \beta} f(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2), \\ M_{14}^t &= -\frac{3m_t^4 \lambda^2 A_t s^2 \Delta_{\tilde{t}_1} \sin \phi_t}{8\pi^2 v^2 \sin^3 \beta \cos \alpha} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2}, \\ M_{23}^t &= -\frac{3m_t^4 \lambda A_t \Delta_{\tilde{t}_1} \Delta_{\tilde{t}_2}}{8\pi^2 v \sin \beta \tan \beta} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} - \frac{3m_t^4 \lambda \cos \beta \Delta_{\tilde{t}_1}}{8\pi^2 v \sin^2 \beta} \frac{\log(m_{\tilde{t}_2}^2/m_{\tilde{t}_1}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)}, \\ M_{24}^t &= \frac{3m_t^4 \lambda A_t^2 s \Delta_{\tilde{t}_2} \sin \phi_t}{8\pi^2 v^2 \sin^3 \beta \cos \alpha} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} + \frac{3m_t^4 \lambda A_t s \sin \phi_t}{8\pi^2 v^2 \sin^3 \beta \cos \alpha} \frac{\log(m_{\tilde{t}_2}^2/m_{\tilde{t}_1}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)}, \end{aligned}$$

$$M_{34}^t = - \frac{3m_t^4 \lambda^2 A_t s \Delta_{\tilde{t}_1} \sin \phi_t}{8\pi^2 v \sin^2 \beta \tan \beta \cos \alpha} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2}, \quad (12)$$

where

$$\begin{aligned} \Delta_{\tilde{t}_1} &= A_t \cos \phi_t - \lambda s \cot \beta, \\ \Delta_{\tilde{t}_2} &= A_t - \lambda s \cot \beta \cos \phi_t, \end{aligned} \quad (13)$$

and the dimensionless function g is defined as

$$g(m_x^2, m_y^2) = \frac{m_y^2 + m_x^2}{m_x^2 - m_y^2} \log \frac{m_y^2}{m_x^2} + 2.$$

Likewise, the matrix elements of M^b as

$$\begin{aligned} M_{11}^b &= \frac{3m_b^4 A_b^2 \Delta_{\tilde{b}_1}^2}{8\pi^2 v^2 \cos^2 \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} + \frac{3m_b^4 A_b \Delta_{\tilde{b}_1}}{4\pi^2 v^2 \cos^2 \beta} \frac{\log(m_{\tilde{b}_2}^2/m_{\tilde{b}_1}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)} \\ &\quad + \frac{3m_b^4}{8\pi^2 v^2 \cos^2 \beta} \log \left(\frac{m_{\tilde{b}_1}^2 m_{\tilde{b}_2}^2}{m_b^4} \right), \\ M_{22}^b &= \frac{3m_b^4 \lambda^2 s^2 \Delta_{\tilde{b}_2}^2}{8\pi^2 v^2 \cos^2 \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2}, \\ M_{33}^b &= \frac{3m_b^4 \lambda^2 \Delta_{\tilde{b}_2}^2}{8\pi^2 \cot^2 \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2}, \\ M_{44}^b &= \frac{3m_b^4 \lambda^2 A_b^2 s^2 \sin^2 \phi_b}{8\pi^2 v^2 \cos^4 \beta \cos^2 \alpha} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2}, \\ M_{12}^b &= - \frac{3m_b^4 \lambda A_b s \Delta_{\tilde{b}_1} \Delta_{\tilde{b}_2}}{8\pi^2 v^2 \cos^2 \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} - \frac{3m_b^4 \lambda s \Delta_{\tilde{b}_2}}{8\pi^2 v^2 \cos^2 \beta} \frac{\log(m_{\tilde{b}_2}^2/m_{\tilde{b}_1}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)}, \\ M_{13}^b &= - \frac{3m_b^4 \lambda A_b \Delta_{\tilde{b}_1} \Delta_{\tilde{b}_2}}{8\pi^2 v \cos \beta \cot \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} - \frac{3m_b^4 \lambda \sin \beta \Delta_{\tilde{b}_2}}{8\pi^2 v \cos^2 \beta} \frac{\log(m_{\tilde{b}_2}^2/m_{\tilde{b}_1}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)}, \\ M_{14}^b &= \frac{3m_b^4 \lambda A_b^2 s \Delta_{\tilde{b}_1} \sin \phi_b}{8\pi^2 v^2 \cos^3 \beta \cos \alpha} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} + \frac{3m_b^4 \lambda A_b s \sin \phi_b}{8\pi^2 v^2 \cos^3 \beta \cos \alpha} \frac{\log(m_{\tilde{b}_2}^2/m_{\tilde{b}_1}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)}, \\ M_{23}^b &= \frac{3m_b^4 \lambda^2 s \Delta_{\tilde{b}_2}^2}{8\pi^2 v \cos \beta \cot \beta} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} - \frac{3m_b^4 \lambda^2 s \tan \beta}{8\pi^2 v \cos \beta} f(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2), \\ M_{24}^b &= - \frac{3m_b^4 \lambda^2 A_b s^2 \Delta_{\tilde{b}_2} \sin \phi_b}{8\pi^2 v^2 \cos^3 \beta \cos \alpha} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2}, \\ M_{34}^b &= - \frac{3m_b^4 \lambda^2 A_b s \Delta_{\tilde{b}_2} \sin \phi_b}{8\pi^2 v \cos^2 \beta \cot \beta \cos \alpha} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2}, \end{aligned} \quad (14)$$

where

$$\begin{aligned} \Delta_{\tilde{b}_1} &= A_b - \lambda s \tan \beta \cos \phi_b, \\ \Delta_{\tilde{b}_2} &= A_b \cos \phi_b - \lambda s \tan \beta, \end{aligned} \quad (15)$$

and the matrix elements of M^k as

$$\begin{aligned}
M_{11}^k &= \frac{3m_k^4 \lambda^2 v^2 \sin^2 \beta \Delta_{\tilde{k}_1}^2}{8\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}, \\
M_{22}^k &= \frac{3m_k^4 \lambda^2 v^2 \cos^2 \beta \Delta_{\tilde{k}_1}^2}{8\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}, \\
M_{33}^k &= \frac{3m_k^4 A_k^2 \Delta_{\tilde{k}_2}^2}{8\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2} + \frac{3m_k^4 A_k \Delta_{\tilde{k}_2}}{4\pi^2 s^2} \frac{\log(m_{\tilde{k}_2}^2/m_{\tilde{k}_1}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)} \\
&\quad + \frac{3m_k^4}{8\pi^2 s^2} \log\left(\frac{m_{\tilde{k}_1}^2 m_{\tilde{k}_2}^2}{m_k^4}\right), \\
M_{44}^k &= \frac{3m_k^4 \lambda^2 A_k^2 v^2 \sin^2 \phi_k}{8\pi^2 s^2 \cos^2 \alpha} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}, \\
M_{12}^k &= \frac{3m_k^4 \lambda^2 v^2 \sin 2\beta \Delta_{\tilde{k}_1}^2}{16\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2} - \frac{3m_k^4 \lambda^2 v^2 \sin 2\beta}{16\pi^2 s^2} f(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2), \\
M_{13}^k &= -\frac{3m_k^4 \lambda A_k v \sin \beta \Delta_{\tilde{k}_1} \Delta_{\tilde{k}_2}}{8\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2} \\
&\quad - \frac{3m_k^4 \lambda v \sin \beta \Delta_{\tilde{k}_1}}{8\pi^2 s^2} \frac{\log(m_{\tilde{k}_2}^2/m_{\tilde{k}_1}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)}, \\
M_{14}^k &= -\frac{3m_k^4 \lambda^2 A_k v^2 \sin \beta \Delta_{\tilde{k}_1} \sin \phi_k}{8\pi^2 s^2 \cos \alpha} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}, \\
M_{23}^k &= -\frac{3m_k^4 \lambda A_k v \cos \beta \Delta_{\tilde{k}_1} \Delta_{\tilde{k}_2}}{8\pi^2 s^2} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2} \\
&\quad - \frac{3m_k^4 \lambda v \cos \beta \Delta_{\tilde{k}_1}}{8\pi^2 s^2} \frac{\log(m_{\tilde{k}_2}^2/m_{\tilde{k}_1}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)}, \\
M_{24}^k &= -\frac{3m_k^4 \lambda^2 A_k v^2 \cos \beta \Delta_{\tilde{k}_1} \sin \phi_k}{8\pi^2 s^2 \cos \alpha} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}, \\
M_{34}^k &= \frac{3m_k^4 \lambda A_k^2 v \Delta_{\tilde{k}_2} \sin \phi_k}{8\pi^2 s^2 \cos \alpha} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2} + \frac{3m_k^4 \lambda A_k v \sin \phi_k}{8\pi^2 s^2 \cos \alpha} \frac{\log(m_{\tilde{k}_2}^2/m_{\tilde{k}_1}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)}, \quad (16)
\end{aligned}$$

where

$$\begin{aligned}
\Delta_{\tilde{k}_1} &= A_k \cos \phi_k - \lambda v \tan \alpha, \\
\Delta_{\tilde{k}_2} &= A_k - \lambda v \tan \alpha \cos \phi_k. \quad (17)
\end{aligned}$$

Notice that the matrix elements M_{i4}^t , M_{i4}^b , and M_{i4}^k ($i = 1 - 3$) are proportional to $\sin \phi_t$, $\sin \phi_b$, and $\sin \phi_k$, respectively. If the three CP phases are not all zero, these matrix elements would not be all zero, and therefore the matrix elements M_{i4} ($i = 1 - 3$) would not be zero, which are responsible for the scalar-pseudoscalar mixing and hence CP violation. In short, these three CP phases in the radiative corrections generate the CP mixing between the scalar and pseudoscalar components, implying that the four neutral Higgs bosons are not states of definite CP parity.

The four physical neutral Higgs bosons at the one-loop level are defined as the eigenstates of the squared mass matrix M , with their squared masses as the eigenvalues of M . Let us denote the physical four neutral Higgs bosons as h_i ($i=1$ to 4) and their squared masses as $m_{h_i}^2$ ($i=1$ to 4). We assume that $m_{h_i}^2 \leq m_{h_j}^2$ for $i < j$. Among them, the upper bound on $m_{h_1}^2$, the squared mass of the lightest neutral Higgs boson, may be obtained by noticing that the smallest eigenvalue of a positive symmetric matrix cannot exceed the smaller eigenvalue of its upper left 2×2 submatrix. Thus, the upper bound on $m_{h_1}^2$ at the one-loop level is given as

$$\begin{aligned}
m_{h_1}^2 \leq & \lambda^2 v^2 \sin^2 2\beta + m_Z^2 \cos^2 2\beta + 2g'^2 v^2 (\tilde{Q}_1 \cos^2 \beta + \tilde{Q}_2 \sin^2 \beta)^2 \\
& + \frac{3m_t^4}{8\pi^2 v^2} \frac{(\lambda s \cot \beta \Delta_{\tilde{t}_1} - A_t \Delta_{\tilde{t}_2})^2}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2) \\
& + \frac{3m_t^4}{4\pi^2 v^2} \frac{(\lambda s \cot \beta \Delta_{\tilde{t}_1} - A_t \Delta_{\tilde{t}_2})}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)} \log \left(\frac{m_{\tilde{t}_2}^2}{m_{\tilde{t}_1}^2} \right) + \frac{3m_t^4}{8\pi^2 v^2} \log \left(\frac{m_{\tilde{t}_1}^2 m_{\tilde{t}_2}^2}{m_t^4} \right) \\
& + \frac{3m_b^4}{8\pi^2 v^2} \frac{(\lambda s \tan \beta \Delta_{\tilde{b}_1} - A_b \Delta_{\tilde{b}_2})^2}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2) \\
& + \frac{3m_b^4}{4\pi^2 v^2} \frac{(\lambda s \tan \beta \Delta_{\tilde{b}_1} - A_b \Delta_{\tilde{b}_2})}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)} \log \left(\frac{m_{\tilde{b}_2}^2}{m_{\tilde{b}_1}^2} \right) + \frac{3m_b^4}{8\pi^2 v^2} \log \left(\frac{m_{\tilde{b}_1}^2 m_{\tilde{b}_2}^2}{m_b^4} \right) \\
& + \frac{3m_k^4 \lambda^2 v^2 \Delta_{k_1}^2}{8\pi^2 s^2} \frac{g(m_{k_1}^2, m_{k_2}^2)}{(m_{k_2}^2 - m_{k_1}^2)^2} - \frac{3m_k^2 \lambda^2 v^2 \sin^2 2\beta}{16\pi^2 s^2} f(m_{k_1}^2, m_{k_2}^2), \quad (18)
\end{aligned}$$

where the first three terms come from the tree-level Higgs potential while the other terms come from the one-loop corrections due to top quark, bottom quark, the exotic quark and their superparticles.

III. NUMERICAL ANALYSIS

For numerical analysis, we assume that the extra $U(1)'$ emerges from the E_6 group, as described in the previous section. To be concrete, we take the ν -model [19], where $U(1)'$ is a mixture of $U(1)_\chi$ and $U(1)_\psi$ with the mixing angle $\theta_E = \tan^{-1} \sqrt{15}$. At the electroweak scale, the analysis of renormalization group equation leads to $g'_1(m_Z) \simeq 0.46$ for $U(1)'$ gauge coupling constant [19]. Then, the $U(1)'$ charges of H_1 , H_2 , and S are given respectively as $\tilde{Q}_1 \approx -0.4910123$, $\tilde{Q}_2 \approx -0.2995571$, and $\tilde{Q}_3 = -(\tilde{Q}_1 + \tilde{Q}_2)$. We set quark masses as $m_t = 175$ GeV, $m_b = 4$ GeV, and $m_k = 700$ GeV. The renormalization scale is set as $\Lambda = 700$ GeV in the one-loop effective potential. We assume that the lighter squarks are larger than top quark mass. For the remaining parameters, we set the ranges for their variations as follows: $100 \leq A_t = A_b = A_k, \bar{m}_A \text{ (GeV)} \leq 2000$, $100 \leq m_{\text{SUSY}} = m_T = m_B = m_K \text{ (GeV)} \leq 1000$, $1500 \leq s \text{ (GeV)} \leq 2000$, $0 \leq \phi_t, \phi_b, \phi_k \leq 2\pi$, $1 < \tan \beta \leq 30$, and $0 < \lambda \leq 0.85$. Note that one need not vary ϕ_0 since it appears always together with A_λ in the definition of $\bar{m}_A$. Also note that the upper bound on λ is chosen by considering that the maximum value of λ is about 0.83 in the analysis of the renormalization group equation. A general comment on s may be that a large s leads

to a heavy Z' in the present model. Moreover, the large s can satisfy the experimental constraint that the mixing angle between Z and Z' should be smaller than $2\text{--}3 \times 10^{-3}$.

We calculate the upper bound on m_{h_1} by varying all the relevant parameters within their allowed ranges for a given $\tan\beta$. The results are shown in Fig. 1, where the upper bound on m_{h_1} is plotted as a function of $\tan\beta$. The solid curve is obtained by considering only the tree-level Higgs potential, whereas the dashed curve is obtained by considering the full one-loop Higgs potential with explicit CP violation. It is observed in the MSSM that the lightest neutral Higgs boson mass increases steadily as $\tan\beta$ increases. In the present model, where the MSSM is incorporated with an extra $U(1)'$, the lightest neutral Higgs boson mass does not increase in accordance with the increase of $\tan\beta$, because of both the Higgs singlet and V_D contributions due to an extra $U(1)'$. One can see that in Fig. 1 the lightest neutral Higgs boson mass becomes maximum at $\tan\beta \sim 1.5$ for both curves.

As we have remarked in the previous section, the scalar-pseudoscalar mixing, which is responsible for CP violation in this model, is triggered by three CP phases, ϕ_t , ϕ_b , and ϕ_k . The size of the CP mixing between the scalar and pseudoscalar Higgs bosons could be regarded as maximal if $\sin\phi_t = \sin\phi_b = \sin\phi_k = 1$, while there would be no CP mixing if $\sin\phi_t = \sin\phi_b = \sin\phi_k = 0$. In order to see a more clear picture about the amount of the CP mixing among the scalar and pseudoscalar Higgs bosons, we introduce ($i = 1, 2, 3$)

$$\omega_i(\phi_k) = \frac{2|M_{i4}|}{|M_{ii}| + |M_{44}|}, \quad (19)$$

which depend implicitly on other parameters as well.

These $\omega_i(\phi_k)$ measure by what amount ϕ_k contributes to the size of CP mixing. In order to express the relative contributions by ϕ_k , we further introduce

$$\Omega_i(\phi_k) = \frac{\omega_i(\phi_k)}{\omega_i(\phi_k = 0)}. \quad (20)$$

By evaluating $\Omega_i(\phi_k)$, we may study the role of the exotic quark sector in the CP violation in the present model.

In Fig. 2, we plot Ω_3 as a function of ϕ_k , where we fix $m_{\text{SUSY}} = \bar{m}_A = A_t = s/2 = 1000$ GeV and $\lambda = 0.3$, and we set for simplicity $\phi_t = \phi_b$. The four curves in Fig. 2 correspond to four different sets of $\phi_t = \phi_b$ and $\tan\beta$, namely, the solid curve is obtained for $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 2$, the dashed curve for $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 20$, the dotted curve for $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 2$, and the dash-dotted curve for $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 20$. It is trivial to notice that $\Omega_3 = 1$ at $\phi_k = 0, \pi$, or 2π . For other values of ϕ_k , we find that Ω_3 may change up to as much as about 5×10^3 for some parameter values. This implies that the effect of ϕ_k upon Ω_3 , or ω_3 , is very significant. In other words, the matrix element $|M_{34}|$ depends critically on the complex phase of the exotic quark sector.

We study the behaviors of Ω_1 and Ω_2 for $0 \leq \phi_k \leq 2\pi$, fixing the values of other parameters as in Fig. 2. Unlike Ω_3 , they do not fluctuate widely against the variation of ϕ_k . Thus, it can be said that the complex phase of the exotic quark sector has a strong influence on the magnitude of the mixing between h_3 and h_4 , whereas it has relatively weak influence on the magnitudes of the mixing between h_1 and h_4 or between h_2 and h_4 .

We calculate the masses of the neutral Higgs bosons for the same parameter setting as in Fig. 2: $m_{\text{SUSY}} = \bar{m}_A = A_t = s/2 = 1000$ GeV, $\lambda = 0.3$, together with four different sets of $\phi_t = \phi_b$ and $\tan\beta$, for $0 \leq \phi_k \leq 2\pi$. We obtain approximately 126, 1000, 1002, and 1043 GeV, respectively, for m_{h_1} , m_{h_2} , m_{h_3} , and m_{h_4} . We find that these values are almost unchanged for $0 \leq \phi_k \leq 2\pi$, and nearly the same within a few GeV for the four different sets of $\phi_t = \phi_b$ and $\tan\beta$.

The mass of the extra gauge boson Z' is also estimated. We find that $m_{Z'}$ is stable at approximately 1039 GeV for the parameter values in Fig. 2. The mixing angle between Z and Z' , denoted as $\alpha_{ZZ'}$, depends on both $\tan\beta$ and s . We find that $\alpha_{ZZ'} = 1.39 \times 10^{-3}$ for $\tan\beta = 2$, and slightly increases for larger values of $\tan\beta$ as 2.93×10^{-3} for $\tan\beta = 20$.

We have studied the dependence of the mixing element between h_3 and h_4 on the complex phase of the exotic quark sector. Meanwhile, one can evaluate effectively the absolute size of the CP violation by calculating the dimensionless parameter

$$\rho(\phi_t, \phi_b, \phi_k) = 4\sqrt{|O_{11}O_{21}O_{31}O_{41}|}, \quad (21)$$

where O_{ij} are the elements of the orthogonal matrix which diagonalizes the squared mass matrix for the neutral Higgs bosons. The range of ρ goes from 0 to 1 since the elements of the orthogonal matrix satisfy the orthogonality condition of $\sum_{j=1}^4 O_{j1}^2 = 1$. If $\rho = 0$, there would be no CP violation, whereas the CP symmetry would be maximally violated if $\rho = 1$. The maximal CP violation that leads to $\rho = 1$ takes place when $O_{11}^2 = O_{21}^2 = O_{31}^2 = O_{41}^2 = 1/4$.

In order to figure out the dependence of ρ on ϕ_k , we introduce

$$\rho_k(\phi_k) = \frac{\rho(\phi_t, \phi_b, \phi_k)}{\rho(\phi_t, \phi_b, \phi_k = 0)} \quad (22)$$

We calculated ρ_k as a function of ϕ_k , for the same parameter setting as Fig. 2. If ρ_k remains at 1, it would imply that ρ_k does not depend on ϕ_k . On the other hand, ϕ_k would contribute more significantly to CP violation if ρ_k moves farther away from 1. The result is shown in Fig. 3. The four curves correspond to the four sets of parameter values. Fig. 3 shows that the fluctuation of ρ_k is larger for $\tan\beta = 20$ than for $\tan\beta = 2$. This implies that ϕ_k play an important role in ρ_k for large $\tan\beta$. One can see that ρ_k fluctuates by up to 20 % for nonzero ϕ_k . Since $\rho(\phi_t, \phi_b, \phi_k)$ is an absolute measure for the CP violation in our model, it is reasonable to expect that the complex phase of the exotic quark sector may change up to 20 % of the CP violation.

IV. SPONTANEOUS CP VIOLATION

In this section, let us examine briefly whether it is possible to realize the spontaneous CP violation in the present model. At initial stage, the Lagrangian density of the model is assumed to be invariant with respect to CP property. After the electroweak symmetry breaking, the CP symmetry may spontaneously be broken by complex phases in the VEVs of the neutral Higgs fields. In the MSSM with an extra $U(1)'$, one can conjecture from the

Georgi-Pais theorem that the spontaneous CP violation leads to a very light pseudoscalar Higgs boson, namely, the axion. It has been addressed that the spontaneous CP violation through radiative corrections can be realized when at the tree level there exist massless Higgs bosons other than Goldstone boson [23]. In the present model, there is no pseudo-Goldstone boson, which is referred to as axion, and thus the spontaneous CP violation cannot occur at the one-loop level.

In order to examine the possibility of spontaneous CP violation scenario at the one-loop level, we can quantitatively analyze the axion mass by using the formulae which are derived for explicit CP violation scenario. In spontaneous CP violation scenario, the one-loop effective potential of the present model may have only one physical CP phase (ϕ_s) arising from the VEV of the Higgs singlet by considering the radiative corrections due to top quark, bottom quark, the exotic quark, and their superpartners. At the one-loop level, we obtain the minimum condition for the vacuum with respect to ϕ_s as

$$A_\lambda = \frac{3m_t^2 A_t}{16\pi^2 v^2 \sin^2 \beta} f(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2) + \frac{3m_b^2 A_b}{16\pi^2 v^2 \cos^2 \beta} f(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2) + \frac{3m_k^2 A_k}{16\pi^2 s^2} f(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2) \quad (23)$$

for $\phi_s \neq 0, \pi$. We obtain the axion mass at the one-loop level as

$$m_a^2 = \frac{3m_t^4 \lambda^2 A_t^2 s^2 \sin^2 \phi_t}{8\pi^2 v^2 \sin^4 \beta \cos^2 \alpha} \frac{g(m_{\tilde{t}_1}^2, m_{\tilde{t}_2}^2)}{(m_{\tilde{t}_2}^2 - m_{\tilde{t}_1}^2)^2} + \frac{3m_b^4 \lambda^2 A_b^2 s^2 \sin^2 \phi_b}{8\pi^2 v^2 \cos^4 \beta \cos^2 \alpha} \frac{g(m_{\tilde{b}_1}^2, m_{\tilde{b}_2}^2)}{(m_{\tilde{b}_2}^2 - m_{\tilde{b}_1}^2)^2} + \frac{3m_k^4 \lambda^2 A_k^2 v^2 \sin^2 \phi_k}{8\pi^2 s^2 \cos^2 \alpha} \frac{g(m_{\tilde{k}_1}^2, m_{\tilde{k}_2}^2)}{(m_{\tilde{k}_2}^2 - m_{\tilde{k}_1}^2)^2}. \quad (24)$$

The massless function g in the above expression is found to have negative or very small values for the entire parameter space we consider, in the case of explicit CP violation scenario. Employing this result from the explicit CP violation scenario, the squared mass of axion is almost always negative for the entire parameter space, which is unacceptable. Therefore, the spontaneous CP violation is not viable for the MSSM with an extra $U(1)'$.

Furthermore, the lightest neutral Higgs boson mass also faces difficulty in the spontaneous CP violation scenario. Calculate the lightest neutral Higgs boson mass from the 4×4 squared mass matrix for neutral Higgs bosons, we find that the squared mass of the lightest Higgs boson has negative values for the above parameter space. In fact, the minimum condition for the vacuum with respect to ϕ_s has a very strong constraint, and thus the vacuum becomes unstable for the full parameter space we consider, in the spontaneous CP violation scenario.

V. CONCLUSIONS

We consider CP violation in the MSSM with an extra $U(1)'$ symmetry, which is assumed to be originated from a string-inspired E_6 model. The charge of the extra $U(1)'$ is defined as $Q' = \cos \theta_E Q_\chi + \sin \theta_E Q_\psi$ with $\theta_E = \tan^{-1} \sqrt{15}$. This model possesses one Higgs singlet and exotic quarks and squarks, in addition to the particle content of the MSSM. In this

model, the CP violation is impossible to occur either explicitly or spontaneously in its tree-level Higgs sector. The mixing between the scalar and pseudoscalar Higgs bosons, which induce the CP violation, may occur via the radiative corrections at the one-loop level. For the radiative corrections, we consider the loop contributions due to top quark, bottom quark, the exotic quark and their superpartners. The radiative corrections contain in general three complex phases originated from the top squark sector, the bottom squark sector, and the exotic squark sector. We pay attention to the complex phase ϕ_k that comes from the exotic squark sector.

We calculate the lightest neutral Higgs boson mass. In explicit CP violation scenario, we find that the mass of the lightest neutral Higgs boson is smaller than about 135 GeV and 165 GeV, respectively, at the tree level and at the one-loop level, for the whole parameter space that we considered.

For four different sets of parameters, we calculate the magnitude of the CP mixing among the scalar and pseudoscalar Higgs bosons for $0 \leq \phi_k \leq 2\pi$. The matrix element, $|M_{34}|$, representing the CP mixing between the heaviest scalar and the pseudoscalar Higgs bosons, is found to depend heavily on the complex phase from the exotic squark sector, ϕ_k . In particular, the magnitude of the CP mixing between the heaviest scalar and the pseudoscalar Higgs bosons is found to change as much as about 20 % for $0 \leq \phi_k \leq 2\pi$ when $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 20$, whereas the CP mixing among the other scalar Higgs bosons and the pseudoscalar Higgs boson is found to be relatively stable against the variation of ϕ_k .

Also, we have investigated the possibility of spontaneous CP violation at the one-loop level in the neutral Higgs sector, where the tree-level Higgs potential possesses CP symmetry. We argue that it is impossible to achieve spontaneous CP violation in the radiatively corrected Higgs potential of this model.

ACKNOWLEDGMENTS

This research is supported by KOSEF through CHEP. The authors would like to acknowledge the support from KISTI (Korea Institute of Science and Technology Information) under "The Strategic Supercomputing Support Program" with Dr. Kihyeon Cho as the technical supporter. The use of the computing system of the Supercomputing Center is also greatly appreciated.

- [1] H.P. Nilles, Phys. Rep. **110**, 1 (1984); J.F. Gunion, H.E. Haber, G.L. Kane, and S. Dawson, *The Higgs Hunters' Guide* (Addison-Wesley Redwood City, CA, 1990).
- [2] S. Weinberg, Phys. Rev. Lett. **37**, 657 (1976).
- [3] A. Pomarol, Phys. Lett. B **287**, 331 (1992); N. Maekawa, Phys. Lett. B **282**, 387 (1992); N. Haba, Phys. Lett. B **398**, 305 (1997).

- [4] O. Lebedev, Eur. Phys. J. C **4**, 363 (1998).
- [5] A. Pilaftsis, Phys. Lett. B **435**, 88 (1998); A. Pilaftsis, Phys. Rev. D **58**, 096010 (1998).
- [6] D.A. Demir, Phys. Rev. D **60**, 055006 (1999); S.Y. Choi, M. Drees, and J.S. Lee, Phys. Lett. B **481**, 57 (2000).
- [7] G.L. Kane and L.T. Wang, Phys. Lett. B **488**, 383 (2000).
- [8] S. Heinemeyer, Eur. Phys. J. C **22**, 521 (2001).
- [9] T. Ibrahim and P. Nath, Phys. Rev. D **63**, 035009 (2001); T. Ibrahim and P. Nath, Phys. Rev. D **66**, 015005 (2002); S.W. Ham, S.K. Oh, E.J. Yoo, C.M. Kim, and D. Son, Phys. Rev. D **68**, 055003 (2002).
- [10] J.S. Lee, A. Pilaftsis, M. Carena, S.Y. Choi, M. Drees, J.R. Ellis, and, C.E.M. Wagner, Comput. Phys. Commun. **156**, 283 (2004).
- [11] J.R. Ellis, J.S. Lee, and A. Pilaftsis, Phys. Rev. D **72**, 095006 (2005).
- [12] A. Pilaftsis and C.E.M. Wagner, Nucl. Phys. **B553**, 3 (1999); M. Carena, J. Ellis, A. Pilaftsis, and C.E.M. Wagner, Nucl. Phys. **B586**, 92 (2000); S.W. Ham, S.K. Oh, E.J. Yoo, and H.K. Lee, J. Phys. G **27**, 1 (2001); M. Carena, J.R. Ellis, A. Pilaftsis, and C.E.M. Wagner, Phys. Lett. B **495**, 155 (2000).
- [13] J.E. Kim and H.P. Nilles, Phys. Lett. B **138**, 150 (1984).
- [14] J. Ellis, J.F. Gunion, H.E. Haber, L. Roszkowski, and F. Zwirner, Phys. Rev. D **39**, 844, (1989).
- [15] J.L. Hewett and T.G. Rizzo, Phys. Rep. **183**, 193 (1989); A. Leike, Phys. Rep. **317**, 143 (1999).
- [16] M. Cvetič and P. Langacker, Phys. Rev. D **54**, 3570 (1996).
- [17] M. Cvetič, D.A. Demir, J.R. Espinosa, L. Everett, and P. Langacker, Phys. Rev. D **54**, 3570 (1996).
- [18] D.A. Demir and N.K. Pak, Phys. Rev. D **57**, 6609 (1998); Y. Daikoku and D. Suematsu, Phys. Rev. D **62**, 095006 (1998); H. Amini, New J. Phys. **5**, 49 (2003).
- [19] S.F. King, S. Moretti, and R. Nevzorov, Phys. Rev. D **73**, 035009 (2006); S.F. King, S. Moretti, and R. Nevzorov, Phys. Lett. B **634**, 278 (2006).
- [20] D.A. Demir and L.L. Everett, Phys. Rev. D **69**, 015008 (2004).
- [21] J. Erler, Nucl. Phys. **B586**, 73 (2000).
- [22] S. Coleman and E. Weinberg, Phys. Rev. D **7**, 1888 (1973).
- [23] H. Georgi and A. Pais, Phys. Rev. D **10**, 1246 (1974).

FIGURE CAPTION

FIG. 1. : The upper bound on the lightest neutral Higgs boson mass as a function of $\tan\beta$, where other relevant parameters are allowed to vary within their respective ranges: $100 \leq A_t = A_b = A_k, \bar{m}_A \text{ (GeV)} \leq 2000$, $100 \leq m_{\text{SUSY}} = m_T = m_B = m_K \text{ (GeV)} \leq 1000$, $1500 \leq s \text{ (GeV)} \leq 2000$, $0 \leq \phi_t, \phi_b, \phi_k \leq 2\pi$, $1 < \tan\beta \leq 30$, and $0 < \lambda \leq 0.85$. The quark masses are set as $m_t = 175 \text{ GeV}$, $m_b = 4 \text{ GeV}$, $m_k = 700 \text{ GeV}$, and the renormalization scale is set as $\Lambda = 700 \text{ GeV}$. The solid and dashed curves are respectively obtained from the tree-level and the one-loop level Higgs potential with explicit CP violation.

FIG. 2. : Plot of Ω_3 as a function of ϕ_k , for four different sets of $\phi_t = \phi_b$ and $\tan\beta$: $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 2$ (solid curve), $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 20$ (dashed curve), $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 2$ (dotted curve), and $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 20$ (dash-dotted curve). The remaining parameters are set as $m_{\text{SUSY}} = \bar{m}_A = A_t = s/2 = 1000 \text{ GeV}$ and $\lambda = 0.3$.

FIG. 3. : Plot of ρ_k as a function of ϕ_k , for the same parameter setting as Fig. 2.




FIG. 1: The upper bound on the lightest neutral Higgs boson mass as a function of $\tan \beta$, where other relevant parameters are allowed to vary within their respective ranges: $100 \leq A_t = A_b = A_k, \bar{m}_A$ (GeV) ≤ 2000 , $100 \leq m_{\text{SUSY}} = m_T = m_B = m_K$ (GeV) ≤ 1000 , $1500 \leq s$ (GeV) ≤ 2000 , $0 \leq \phi_t, \phi_b, \phi_k \leq 2\pi$, $1 < \tan \beta \leq 30$, and $0 < \lambda \leq 0.85$. The quark masses are set as $m_t = 175$ GeV, $m_b = 4$ GeV, $m_k = 700$ GeV, and the renormalization scale is set as $\Lambda = 700$ GeV. The solid and dashed curves are respectively obtained from the tree-level and the one-loop level Higgs potential with explicit CP violation.




FIG. 2: Plot of Ω_3 as a function of ϕ_k , for four different sets of $\phi_t = \phi_b$ and $\tan\beta$: $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 2$ (solid curve), $\phi_t = \phi_b = \pi/6$ and $\tan\beta = 20$ (dashed curve), $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 2$ (dotted curve), and $\phi_t = \phi_b = \pi/3$ and $\tan\beta = 20$ (dash-dotted curve). The remaining parameters are set as $m_{\text{SUSY}} = \bar{m}_A = A_t = s/2 = 1000$ GeV and $\lambda = 0.3$.




FIG. 3: Plot of ρ_k as a function of ϕ_k , for the same parameter setting as Fig. 2.